

AD-A048 841

ILLINOIS UNIV AT URBANA-CHAMPAIGN DEPT OF CHEMISTRY
BIOLUMINESCENCE OF THE FIREFLY. KEY STEPS IN THE FORMATION OF T--ETC(U)
OCT 77 J KOO, S P SCHMIDT, G B SCHUSTER

N00014-76-C-0745

NL

UNCLASSIFIED

TR-06

1 OF 1

ADAO48841



END

DATE
FILMED

2 - 78

DDC

DEC 29 1977

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS
BEFORE COMPLETING FORM

1. REPORT NUMBER D N0014-76-C-0745-06	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Bioluminescence of the Firefly. Key Steps in the Formation of the Electronically Excited State for Model Systems.		5. TYPE OF REPORT & PERIOD COVERED Technical
7. AUTHOR(s) Gary B. Schuster Ja-Young Koo Steven P. Schmidt		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry University of Illinois Urbana, Illinois 61801		8. CONTRACT OR GRANT NUMBER(s) N0014-76-C-0745
11. CONTROLLING OFFICE NAME AND ADDRESS Chemistry Program, Materials Science Division Office of Naval Research, 800 N. Quincy St., Arlington, Va 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-051-616
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE October 14, 1977
		13. NUMBER OF PAGES 13
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE

15. DISTRIBUTION STATEMENT (of this Report)

Approved for Public Release, Distribution Unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

18. SUPPLEMENTARY NOTES

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

chemiluminescence
bioluminescence
dioxetane
electron transfer

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

The chemical mechanism for formation of electronically excited state molecules from the thermal reaction of dimethyldioxetanone was probed. Light production in the presence of certain easily oxidized aromatic hydrocarbons was found not to conform to the classical mechanistic schemes for chemiexcitation. Detailed investigation of the dioxetanone system revealed light formation by the recently discovered chemically initiated systems. In particular, the key high energy molecule involved in firefly luminescence, which has been identified as a dioxetanone, is postulated to form excited

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 63 IS OBSOLETE
S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DDC FILE COPY

AD A 048841

DDC
RECEIVED
JAN 18 1978
A

20. Abstract

states as a result of intramolecular electron transfer from the phenoxythiazole moiety to the dioxetanone. Subsequent rapid decarboxylation results in direct formation of an excited singlet state of the emitting amide.

RECEIVED
JAN 18 1978
A

158840A DA
JUL 31 1978

OFFICE OF NAVAL RESEARCH

Contract ~~NO014-76-C-0745~~

Task No. NR-051-616

TECHNICAL REPORT NO. ~~NO014-76-C-0745-06~~

Bioluminescence of the Firefly. Key Steps in the
Formation of the Electronically Excited State for Model Systems.

by Technical rept.,

Ja-Young/Koo, Steven P./Schmidt, Gary B./Schuster

Prepared for Publication

in

Proceedings of the National

Academy of Science U.S.A.

Dept of Chemistry
School of Chemical Sciences
University of Illinois
Urbana, Illinois 61801

11/14 Oct 1977

12/17 P.

Reproduction in whole or in part is permitted for
any purpose of the United States Government

Approved for Public Release; Distribution Unlimited.

ACCESSION for	
NMS	White Section <input checked="" type="checkbox"/>
DIC	Dist Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

408 087

mt



Classification: Chemistry

Bioluminescence of the Firefly. Key Steps in the
Formation of the Electronically Excited State for Model Systems.
(chemiluminescence / electron transfer / dioxetane / fluorescence)

Ja-Young Koo, Steven P. Schmidt, and Gary B. Schuster*

Department of Chemistry
Roger Adams Laboratory
University of Illinois
Urbana, Illinois 61801

*To whom reprint requests should be addressed. Fellow of the Alfred P. Sloan Foundation.


Abstract:  The chemical mechanism for formation of electronically excited state molecules from the thermal reaction of dimethyldioxetanone was probed. Light production in the presence of certain easily oxidized aromatic hydrocarbons was found not to conform to the classical mechanistic schemes for chemiexcitation. Detailed investigation of the dioxetanone system revealed light formation by the recently discovered chemically initiated electron exchange process. This result is extrapolated to bioluminescent systems. In particular, the key high energy molecule involved in firefly luminescence, which has been identified as a dioxetanone, is postulated to form excited states as a result of intramolecular electron transfer from the phenoxythiazole moiety to the dioxetanone. Subsequent rapid decarboxylation results in direct formation of an excited singlet state of the emitting amide. 

Bioluminescent organisms are widely distributed throughout terrestrial and aquatic environments. Although the biological purpose of luminescence varies from species to species, the chemical mechanism for generation of the electronically excited state, which subsequently emits light, appears to be general in a wide variety of organisms. In nearly all of the bioluminescent processes that have been investigated, high-energy cyclic peroxide molecules are implicated as providing the energy necessary for excited state generation (1). In the study of bioluminescent mechanisms the central concerns have been: i) identification of the molecule which has an energy content sufficient to permit excited state generation, ii) characterization of the emitting species, and iii) identification of the molecular process that converts the high-energy content molecule to an electronically excited state.

Our recent investigations of chemiluminescence have led to the discovery of a new general mechanism of excited state formation identified as chemically initiated electron exchange luminescence (2). Studies of exergonic chemical reactions that model bioluminescent systems now permit us to suggest that this mechanism is operative in the formation of electronically excited states in living organisms. The light-forming reaction of the North American Firefly (Photinus pyralis) will serve as the prototypical case. The conclusions reached from this system are readily extended to other bioluminescent reactions.

Many excellent studies of bioluminescence from the firefly have led to the characterization of the enzyme / substrate system involved in the light generation step (3). In summary, the substrate luciferin has been identified as L and independently synthesized. Reaction of L with oxygen in the presence of the enzyme luciferase generates a high energy content molecule which has been identified as the dioxetanone (L_2) by ^{18}O labeling studies (4). In order to produce bioluminescence, L_2 loses the elements of carbon dioxide and forms

an electronically excited singlet state of the observed emitting amide $\tilde{3}$, with very high efficiency. It is the mechanism of this last step, the chemiexcitation step, which has not been previously understood and to which we will direct our attention.



The cyclic peroxidic molecules dioxetane (4) and dioxetanones (5) have been investigated in a number of elegant synthetic and mechanistic studies (5). The observation that the thermal reactions of these high energy compounds lead to the formation of electronically excited carbonyl compounds is taken as strong circumstantial evidence for the involvement of a similar intermediate in the bioluminescence of the firefly. The major difficulty with this interpretation, however, has been the fact that simply substituted



dioxetanes and dioxetanones give rise predominantly to the essentially non-luminescent triplet excited state of the carbonyl compound. On the other hand, the natural bioluminescent systems form the emitting excited singlet state with efficiencies approximating 100%. Application of the concepts of chemically initiated electron exchange luminescence resolves this apparent incongruity.

Experimental

General. Chemiluminescence was detected by the photon counting technique using an EMI 9813 photomultiplier tube. Spectral resolution was achieved with a Jarrel-Ash / Ebert 0.25 m grating monochromator. Emission intensities were corrected for photocathode response using the manufacturer's

spectral sensitivity data and the centers of gravity for the emission spectra. Temperature control was maintained to within $\pm 0.05^\circ$ by means of an external temperature bath. All solvents were purified by passing them through a column of activated alumina followed by distillation. Rubrene, naphthacene, perylene, and 9,10-diphenylanthracene were purchased from Aldrich Chemical Company and purified by chromatography on alumina and recrystallization before use.

Dimethyldioxetaneone (8) was prepared according to the literature procedure of Adam and coworkers (6). This compound was found to be remarkably sensitive to minute quantities of impurities. Extreme care must be used in purification of solvents and additives to suppress the impurity catalyzed decomposition pathways.

Chemiluminescence of 8 in the Presence of Aromatic Hydrocarbons.
Solutions of dioxetaneone 8 in CH_2Cl_2 (typical concentration 10^{-6} - 10^{-5} M) and the appropriate aromatic hydrocarbon (typical concentration 10^{-5} - 10^{-4} M) were prepared. The solutions were deoxygenated by purging with purified, filtered Argon for 3 min at 0° . The chemiluminescent intensity was measured by integrating the light emission at the maximum wavelength for the aromatic hydrocarbon fluorescence.


Results and Discussion

Conventional Mechanisms for Chemiluminescence. Previous studies of chemiluminescent phenomena have led to the evolution of two broad classes of mechanisms for chemiexcitation. In the first, thermal rearrangement of a high-energy content molecule, reacting directly or through an intermediate, generates an electronically excited state of a product molecule. Emission of a photon of light from the excited molecule results in observable direct chemiluminescence. Alternatively, the excited product molecule may enter into a bimolecular energy transfer reaction with a suitable energy acceptor.

This process results in formation of the electronically excited acceptor which, in a subsequent step, emits a photon of light, producing indirect chemiluminescence. This sequence is shown schematically in Figure 1.

The second general mechanism for chemiexcitation that has been well documented is electrogenerated chemiluminescence (7). In this approach a radical cation, usually generated by electrochemical oxidation at a suitable anode, and a radical anion, typically the result of reduction at a cathode, form a diffusive encounter pair. Charge annihilation in the encounter ion pair results in the generation of an electronically excited state. Subsequent emission of light from this excited state, or one derived from it, results in detectable luminescence. This general pathway is shown schematically in Figure 2.

Chemically Initiated Electron Exchange Luminescence. In the course of our studies of the thermal reactions of high-energy content molecules, we established that the chemiluminescence from diphenoyl peroxide (6) did not conform to either of the conventional mechanisms for chemical light formation. The thermolytic conversion of peroxide 6 to benzocoumarin (7), in ca. 70% yield occurs with no significant generation of an electronically excited state of 7. However, it was found that inclusion of any one of several easily oxidized aromatic hydrocarbons in the reaction solution led to efficient



chemiexcitation of the hydrocarbon and bright light emission. Moreover, it was demonstrated that the aromatic hydrocarbon exerted a powerful catalytic effect on the decomposition of diphenoyl peroxide. In addition, it was shown that it is this catalytic path that gives rise to the chemiluminescence.

Significantly, the rate at which the aromatic hydrocarbon catalyzes chemiexcitation, and the actual yield of electronically excited states, correlates with the one-electron oxidation potential of the aromatic hydrocarbon, Figure 3.

Thus, we have identified the initiating step of this reaction sequence as an activated electron transfer from the aromatic hydrocarbon to diphenoyl peroxide.

Subsequent rapid decarboxylation and ring closure generates the radical anion of benzocoumarin within the same solvent cage as the radical cation of the aromatic hydrocarbon. Charge annihilation of these species then generates the luminescing excited state of the aromatic hydrocarbon. This reaction sequence is detailed in Figure 4. Further convincing evidence that this reaction proceeds through a charge annihilation involving benzocoumarin radical anion is derived from the observation of exciplex emission

when triphenylamine is the electron donor. The chemiluminescent exciplex emission in this case is identical in all respects to the emission generated by photoexcitation of mixtures of benzocoumarin and triphenylamine. We have designated this sequence of reactions leading to light emission chemically initiated electron exchange luminescence (CIEEL) (2).


We have demonstrated the generality of chemiexcitation by the CIEEL mechanism in a number of different systems. Of most relevance to the present discussion of bioluminescent mechanisms is the dioxetanone system. Initial investigation of dioxetanone thermolysis led to the observation that unimolecular decomposition generates triplet excited carbonyl compounds in at least twenty times higher yield than the singlet excited carbonyl compound (8). In a later report (9), Adam noted that the overall yield of light was greater by a factor of twenty when rubrene was included in the reaction solution than when 9,10-diphenylanthracene (DPA) was the additive. This result was interpreted as being due to a triplet-triplet annihilation reaction. However, triplet-triplet annihilation as the cause of the increased light yield is inconsistent on

theoretical grounds and has recently been shown experimentally not to be possible under the chemiluminescent conditions reported (10).

We have investigated the chemiluminescent reaction of dimethyldioxetanone (8). Our results show that the rate of reaction of dioxetanone 8 is dependent upon the presence, concentration, and nature of easily oxidized aromatic hydrocarbons. Also, most significantly, for the series of aromatic hydrocarbons, DPA, perylene, 9,10-diphenylethynylanthracene and rubrene; the total light yield is directly correlated with the oxidation potential of the hydrocarbon. These observations implicate CIEEL as the major light producing reaction pathway for dioxetanones. If simple electronic energy transfer from the initially formed singlet excited state of acetone to the aromatic hydrocarbon was responsible for light production, then all of the aromatic hydrocarbons, when corrected for fluorescence efficiency, should have generated the same photon yield.

Bioluminescence by Chemically Initiated Electron Exchange Luminescence.
In considering the options available to the high energy bioluminescent intermediate, dioxetanone 3, for chemiexcitation, it is immediately clear that this molecule is ideally constituted for excited state production by intramolecular chemically initiated electron exchange. In particular, electron transfer from the easily oxidized polycyclic heterocycle portion of the molecule to the high energy dioxetanone moiety followed by rapid decarboxylation will generate a charge transfer resonance structure of the excited singlet state of the observed emitter. The intramolecularity of this process prejudices the reaction toward the formation of excited states of singlet multiplicity, as is observed in the living system. This reaction sequence is outlined in equation 3.

Consistent with this mechanistic postulate, it has been observed that methylation of the phenolic oxygen of firefly luciferin makes the system



non-bioluminescent although the methylated ketone itself is an efficient fluorescer (11). This is the expected result if intramolecular CIEEL is responsible for light production. The methylated substrate is anticipated to be much more difficult to oxidize than the phenol anion (12).

In summary, we have demonstrated that our recently discovered CIEEL mechanism for chemical light formation operates in the dioxetanone system. The dioxetanone is a reasonable model for firefly bioluminescence. Application of CIEEL to these bioluminescent systems results in an explanation for the observed high singlet yield and the effect of methylation. Moreover, chemiexcitation by this path is quite efficient biologically. Nature has built both an effective electron donor and a high energy electron acceptor, as well as an efficient fluorescer, into the same molecule. Certainly this is an effective way to insure high yield light formation. Experiments directed toward the detailed elucidation of the bioluminescent pathway are underway in our laboratory.


Acknowledgment

This work was supported in part by the Office of Naval Research and in part by a grant from the Research Corporation.

References

1. McCapra, F. (1976) "Chemical Mechanisms in Bioluminescence", Accounts Chem. Res., 9, 201-208.
2. Koo, J-Y. and Schuster, G. B. (1977) "Chemically Initiated Electron Exchange Luminescence. A New Chemiluminescent Reaction Path for Organic Peroxides", J. Am. Chem. Soc., 99, 6107-6109.
3. Cormier, M. J., Hercules, D. M. and Lee, J. Ed. (1973) "Chemiluminescence and Bioluminescence", Plenum Press, New York, NY.
4. White, E. H., Maino, J. D., and Umbreit, M. (1975) "On The Mechanism of Firefly Luciferin Luminescence", J. Am. Chem. Soc., 97, 198-200.
5. Wilson, T. (1976), Int. Rev. Sci.: Phys. Chem., 9, 265-311.
6. Adam, W., and Liu, J=C., (1972) "An α -Peroxy Lactone. Synthesis and Chemiluminescence". J. Am. Chem. Soc., 94, 2894-2895.
7. Faulkner, L. R., (1976), Int. Rev. Sci.: Phys. Chem. Ser. Two, 9, 213-264.
8. Turro, N. J., Lechtken, P., Schuster, G., Orell, J., Steinmetzer, H-C., and Adam, W., (1974), "Indirect Chemiluminescence by 1,2-Dioxetanes". J. Am. Chem. Soc., 96, 1627-1629.
9. Adam, W., Simpson, G. A., and Yany, F., (1974) "Mechanism of Direct and Rubrene Enhanced Chemiluminescence During α -Peroxylactone Decarboxylation", J. Phys. Chem., 78, 2559-2569.
10. Liu, D. K. K., and Faulkner, L. R., (1977), "P-Type Delayed Fluorescence from Rubrene", J. Am. Chem. Soc., 99, 4594-4599.
11. McCapra, F., (1970), "Chemiluminescence of Organic Compounds", Pure Appl. Chem., 24, 611-629.
12. Mann, C. K., Barnes, K. K., (1970), "Electric Chemical Reactions in Nonaqueous Systems", Marcel Dekker, New York, NY.

Conventional Chemiluminescence of Organic Molecules




Examples:

a) 1,2-Dioxetanes




b) Luminal




c) Dewar Benzene





Electrogenerated Chemiluminescence (ECL)



MO - Picture







ONR TECHNICAL REPORT DISTRIBUTION LIST

<u>No. Copies</u>	<u>No. Copies</u>
Office of Naval Research ² Arlington, Virginia 22217 Attn: Code 472 2	Commander Naval Undersea Research & Development Center San Diego, California 92132 Attn: Technical Library, Code 133 1
Office of Naval Research Arlington, Virginia 22217 Attn: Code 102IP 6	Naval Weapons Center China Lake, California 93555 Attn: Head, Chemistry Division 1
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. George Sandoz 1	Naval Civil Engineering Laboratory Port Hueneme, California 93041 Attn: Mr. W. S. Haynes 1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept. 1	Professor O. Heinz Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940 1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus 1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380 1
ONR Branch Office 700 Market Street, Rm. 447 San Francisco, California 94102 Attn: Dr. P. A. Miller 1	Dr. M. A. El-Sayed University of California Department of Chemistry Los Angeles, California 90024 1
ONR Branch Office 405 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles 1	Dr. M. W. Windsor Washington State University Department of Chemistry Pullman, Washington 99163 1
Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Library, Code 2029 (ONRL) 6 Technical Info. Div. 1 Code 6100, 6170 1	Dr. E. R. Bernstein Colorado State University Department of Chemistry Fort Collins, Colorado 80521 1
The Asst. Secretary of the Navy (R&D) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350 1	Dr. C. A. Heller Naval Weapons Center Code 6059 China Lake, California 93555 1
Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser) 1	Dr. G. Jones, II Boston University Department of Chemistry Boston, Massachusetts 02215 1
Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314 12	Dr. M. H. Chisholm Chemistry Department Princeton, New Jersey 08540 1
U.S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina Attn: CRD-AA-IP 1	Dr. J. R. MacDonald Code 6110 Chemistry Division Naval Research Laboratory Washington, D.C. 20375 1

ONR TECHNICAL REPORT DISTRIBUTION LIST

No. Copies

Dr. G. B. Schuster
Chemistry Department
University of Illinois
Urbana, Illinois 61801 1

Dr. E. M. Eyring
University of Utah
Department of Chemistry
Salt Lake City, Utah 1

Dr. A. Adamson
University of Southern California
Department of Chemistry
Los Angeles, California 90007 1

Dr. M. S. Wrighton
Massachusetts Institute of Technology
Department of Chemistry
Cambridge, Massachusetts 02139 1